

A classification of explosions in dimension one

E. Sander - J.A. Yorke

Preprint February 2, 2008

Abstract

A discontinuous change in the size of an attractor is the most easily observed type of global bifurcation. More generally, an *explosion* is a discontinuous change in the set of recurrent points. An explosion often results from heteroclinic and homoclinic tangency bifurcations. Newhouse and Palis conjectured in 1976 that planar explosions are generically the result of either tangency or saddle node bifurcations. In this paper, we prove this conjecture for one-dimensional maps. Furthermore, we give a full classification for all possible tangency bifurcations and whether they lead to explosions.

1 Introduction

For continuously varying one-parameter families of iterated maps in R^n , discontinuous changes in the size of an attractor are the most easily observed type of global bifurcations, including changes in the basin boundary (metamorphosis). These changes can occur as the result of a change in stability of the recurrent set. A more general situation occurs when discontinuous changes in attractors occur as the result of a discontinuous change in the size of the recurrent set itself. Such a global bifurcation is called an **explosion**. For the last several years, we have been studying explosions and their properties, including a classification of explosions at heteroclinic tangencies for planar diffeomorphisms [1], and more recently a numerical study of the statistical properties of a certain kind of explosions that occur in dimension three and higher [2]. Our research, as well as that of many others [14, 20], has been guided by a 1976 conjecture of Newhouse and Palis [18]. (See also the re-statements in [10, 19]). For over thirty years, this conjecture has managed to

elude proof. The conjecture says that for generic planar diffeomorphisms, all explosions occur through the following two local bifurcations: saddle-node bifurcations and homoclinic bifurcations. In this paper, we prove that the conjecture is true for smooth interval maps. In addition, we are able to give a full classification of which types of tangencies give rise to explosions. These are important and new result in themselves. They are for example in contrast with a recent result of Horita, Muniz, and Sabini [14], showing that in a probabilistic sense, the Newhouse Palis conjecture is not true for circle maps. In a broader sense, we are hopeful that the insight gained from the one-dimensional case will give rise to insights leading to the proof of the planar case of the Newhouse Palis conjecture.

It is clear that an isolated saddle node bifurcation for either a fixed point or periodic orbit gives rise to a local explosion, since new periodic points appear. However, in many cases in both one and higher dimensions, a saddle node bifurcation also gives rise to a global bifurcation. The set of recurrent points changes discontinuously as the parameter is varied at points not contained in the saddle node periodic orbit. For example, the period three window in the chaotic attractor for the logistic map corresponds to an explosion: For parameter values less than a bifurcation value, there is a global attractor which comprises the full recurrent set. It consists of an interval. After the bifurcation parameter, the global attractor consists only of a period three orbit, and the full recurrent set is a nowhere dense Cantor set within the attractor interval. An explosion occurs at the bifurcation parameter, and it is due to a saddle-node bifurcation at a point inside the global attractor. Saddle node bifurcations on invariant circles are another well-studied examples of this phenomenon.

In all dimensions, explosions due to a homoclinic or heteroclinic bifurcation essentially occur due to the creation of a homoclinic tangle when stable and unstable manifolds intersect transversally. In one and two dimensions, the transition between no intersection and transverse intersections involve tangencies between the stable and unstable manifolds of fixed or periodic points (*cf.* [10]). For example, the Hénon map and Ikeda map contain well studied examples of explosions which are a result of homoclinic bifurcations. In higher dimensions, such bifurcations can occur without tangencies [13]. Such a bifurcation leads to unstable dimension variability [2].

In our previous work for planar maps, we gave a precise classification for which types of tangencies for heteroclinic cycles will result in explosions. We called this class of cycles **crossing cycles**, because the different stable and

unstable manifolds involved in the cycle lie *across* the tangency from each other. We show here that the same results hold for interval maps. Our main results are as follows:

Theorem 1 (Explosions at tangencies). *For generic one-parameter families of smooth maps of the interval with homoclinic orbits or heteroclinic cycles (hypotheses H1-6), explosions occur if and only if there is an isolated crossing orbit.*

Theorem 2 (General explosion classification). *Explosions within generic one-parameter families of smooth maps of the interval (hypotheses H1-3) are the result of either a tangency between stable and unstable manifolds of fixed or periodic points or a saddle node bifurcation of a fixed or periodic point.*

The paper proceeds as follows: In Section 2, we give basic definitions of explosions and homoclinic tangencies. We are considering the particular recurrence class of chain recurrent points, which are defined in this section as well. In Section 3, we prove the Explosions at tangencies theorem. In Section 4, we prove the General explosion classification theorem. Our results rely on a very sophisticated and well-developed theory for the dynamics of interval maps. We have briefly stated the necessary results in the course of the proof.

2 Basic definitions

We now give some formal definitions of concepts described in the introduction. Let $f : (I \times J) \subset (R \times R) \rightarrow R$ be a smooth one-parameter family of maps. We exchangeably write two notations: $f(x, \lambda) = f_\lambda(x)$. For the definition of an explosion it is more natural to use the concept of chain recurrence rather than recurrence. The relationship between chain recurrence and other types of recurrence is discussed in [10].

Definition 1. *For an iterated function g , there is an ϵ -chain from x to y when there is a finite sequence $(z_0, z_1, \dots, z_N)$ such that $z_0 = x$, $z_N = y$, and $d(g(z_{n-1}), z_n) < \epsilon$ for all n .*

*If there is an ϵ -chain from x to itself for every $\epsilon > 0$ (where $N > 0$), then x is said to be **chain recurrent** [11, 12]. The **chain recurrent set** is the set of all chain recurrent points. For a one-parameter family f_λ , we say (x, λ) is chain recurrent if x is chain recurrent for f_λ .*

If for every $\epsilon > 0$, there is an ϵ -chain from x to y and an ϵ -chain from y to x , then x and y are said to be in the same **chain component** of the chain recurrent set.

The chain recurrent set and the chain components are invariant under forward iteration. The following definition is for an explosion bifurcation in the chain recurrent set. Such a definition can be formulated for the non-wandering set as well.

Definition 2 (Chain explosions). A **chain explosion point** (x, λ_0) is a point such that x is chain recurrent for f_{λ_0} , but there is a neighborhood N of x such that on one side of λ_0 (i.e. either for all $\lambda < \lambda_0$ or for all $\lambda > \lambda_0$), no point in N is chain recurrent for f_λ . (All explosion points in this paper are chain explosions, so we sometimes drop the qualifier chain.)

Remark 1. In order to show that (y, λ) is not chain recurrent, it is sufficient to show that y is in the closure of the hyperbolic periodic orbits for f_λ .

Remark 2. In the above definition, at f_{λ_0} , x is not necessarily an isolated point of the chain recurrent set. For example, at a saddle node bifurcation on an invariant circle, the chain recurrent set consists of two fixed points prior to bifurcation and the whole circle at and in many cases after bifurcation.

The chain recurrent set is not invariant under backwards iteration of a noninvertible map. Thus explosion points are not preserved under iteration, forward or backward. The following lemma states what is guaranteed by the fact that chain recurrence is preserved under forward iteration.

Lemma 1. Let (x, λ_0) be a chain explosion point for f . Specifically, there exists $\delta > 0$ such that there is no chain recurrent point in $B_\delta(x)$ for all $\lambda < \lambda_0$, but x is chain recurrent at λ_0 . Then $f(x)$ is chain recurrent at λ_0 , but $f(x)$ may also be chain recurrent for $\lambda < \lambda_0$. In contrast, if x_{-1} is a preimage of x , then there is a $\delta_{-1} > 0$ such that no point in $B_{\delta_{-1}}(x_{-1})$ is chain recurrent for all $\lambda < \lambda_0$. Note that x_{-1} may not be chain recurrent at λ_0 .

We now give definitions of homoclinic and heteroclinic points. Note that for a diffeomorphism, homoclinic and heteroclinic orbits require the existence of saddle points with stable and unstable manifolds of dimension at




Figure 1: A one-dimensional map with a repelling fixed point x_0 where $f'(x_0) > 0$. The point y is a homoclinic point, since $f^k(y) = x_0$ (for $k = 1$), and there exists a sequence of successive preimages of y converging to x_0 .

least one. Thus they can only occur in dimension two or greater. However, for noninvertible maps, it is possible to have fixed or periodic points with one-dimensional unstable manifolds and a zero-dimensional stable manifolds. Marotto terms such points snap-back repellers [17]. It is not possible to reverse these stable and unstable manifold dimensions; the existence of a homoclinic orbit to an attracting fixed point requires a multivalued map [21]. The following definition of homoclinic points for interval maps is depicted in Figure 1.

Definition 3 (Homoclinic points). *Let $f : R \rightarrow R$ be a smooth function with a repelling fixed point x_0 . Let y be a point in the unstable manifold of x_0 and an integer $K > 0$ such that $f^K(y) = x_0$. Then y is a homoclinic point to x_0 . If x_0 is periodic with least period m , then the same definition applies by replacing f with f^m .*

Remark 3. *If p is a hyperbolic fixed point, and the forward limit set of a point z in the unstable manifold of p includes p , then there are two possibilities: (1) a finite iterate of z is equal to p , (2) the limit set of z contains points other than p . In this second case, it is not possible for the limit set of z to consist of p alone. Therefore z is not referred to as a homoclinic point, since its limit set is larger than just p . This case is considered in later sections of this paper.*




Figure 2: A one-dimensional map with a repelling fixed point x_0 with $f'(x_0) > 0$. The point w is a homoclinic tangency point. For the particular tangency depicted here, w is contained in a *non-crossing orbit* (Definition 7), and thus by Theorem 6 w is not an explosion point.

For diffeomorphisms, all orbits through homoclinic points are homoclinic orbits. For one-dimensional maps, there may be many non-homoclinic orbits through a homoclinic point.

Definition 4 (Homoclinic orbits). *Let f, x_0 , and y be as in the above definition of a homoclinic point. An orbit $(z_{-k})_{k=0}^{\infty}$ is a homoclinic orbit through y if the following conditions are satisfied: $z_0 = x_0$, $z_{-K} = y$ for some K , and for all $k \in \mathbb{N}$, $f(z_{-k}) = z_{-k+1}$, and $\lim_{k \rightarrow \infty} z_{-k} = x_0$.*

Since the stable manifold of a homoclinic point is zero-dimensional, a homoclinic tangency is a tangency of the graph of the map at a homoclinic point. Homoclinic tangencies are depicted in Figures 2 and 3.

Definition 5 (Homoclinic tangencies). *Let f, x_0, y , and $(z_{-k})_{k=0}^{\infty}$ be as in the above definition of a homoclinic orbit. The point $w = z_{-L}$ is a homoclinic tangency point if the graph of f is tangent to the horizontal line at w .*

3 Explosions at homoclinic tangencies

This section classifies explosions occurring via homoclinic tangencies. The results are stated for fixed points, but the same results hold for homoclinic




Figure 3: In a homoclinic orbit as in Figure 2, the tangency point w does not need to be the immediate preimage of the fixed point. Here, w is the second preimage of x_0 .

orbits for periodic points with least period m if f is replaced by $g = f^m$. Throughout this section, we make the following hypotheses. The first hypothesis is a smoothness assumption. The second and third hypotheses are generic assumptions for one-parameter families. The fourth is a notational convention for the existence of a homoclinic orbit. The fifth hypothesis is generic for one-parameter families containing a homoclinic orbit.

H1 $f : (I \times J) \subset (R \times R) \mapsto I$ is a C^1 smooth family of C^2 interval maps. For a fixed parameter λ_0 , denote $f_0 := f_{\lambda_0}$.

H2 Assume H1. Assume that there are no intervals on which f_0 is constant.

H3 Assume H1. Assume that for f_0 , there is at most one of the following:

- (a) A non-hyperbolic fixed point or periodic orbit. We assume generic behavior as a parameter is varied. That is, non-hyperbolic point is a codimension one period doubling or saddle-node bifurcation.
- (b) One critical point which comprises a tangency between stable and unstable manifolds of fixed points or periodic orbits. We assume generic behaviour as a parameter is varied. That is, with variation of parameter, the critical point moves from above to below (or below to above) the placement of the periodic point.

- H4** Assume H1, and that for each λ , x_λ is a repelling fixed point for f_λ . Denote $x_0 := x_{\lambda_0}$. Assume that for f_0 , y is homoclinic to x_0 .
- H5** Assume H1 and H4. At $\lambda = \lambda_0$, the homoclinic orbit containing y contains only one critical point.

The following theorem says that if at a repelling periodic point a map has negative derivative then no homoclinic points are explosion points.

Theorem 3 (No explosions for orbits with negative derivatives). *Assume that f_λ is a family of maps satisfying H1-5, and $f'_{\lambda_0}(x_0) < 0$, then (y, λ_0) is not an explosion point.*

Proof. We show that y is contained in the closure of the hyperbolic periodic points of f_{λ_0} .

If y is contained in a homoclinic orbit without a tangency, then the homoclinic orbit is preserved under perturbation, which automatically implies that (y, λ_0) is not an explosion point. Thus we assume that y is contained in a homoclinic orbit containing a homoclinic tangency point w . As before, denote this orbit by $(z_{-k})_{k=0}^\infty$, where $z_0 = x_0$, $\lim_{k \rightarrow \infty} z_{-k} = x_0$, and K and L are such that $y = z_{-K}$ and $w = z_{-L}$.

Fix a neighborhood U of y . By H5, there exists a sequence of neighborhoods $U_k \ni z_{-k}$, such that: (i) $U_K \subset U$, (ii) for $k \geq L$, $f_0(U_{k+1}) = U_k$ and the map is injective, and (iii) for $0 \leq k < L$, $U_{L-k} = f_0^k(U_L)$. Let $V_1 = f_0^L(U_L \setminus w) = U_0 \setminus x_0$. By H2, V_1 is an interval on one side of x_0 . By assumption $f'_0(x_0) < 0$, so if the $\{U_k\}$ are sufficiently small, $V_2 = f_0^{L+1}(U \setminus w)$ is an interval on the other side of x_0 . That is, $V_1 \cup V_2 \cup \{x_0\}$ is a neighborhood of x_0 .

For any sufficiently large J , $U_J \subset V_i$, where $i \in \{1, 2\}$. In addition, $f^M(U_J) = V_i$, where $M = J$ or $M = J + 1$. Thus for $0 \leq k \leq J$, U_k contains a periodic point. Thus there is a periodic point in U . Since U was arbitrary, there are periodic points p_j with period n_j such that $\lim_{j \rightarrow \infty} p_j = y$ and $\lim_{j \rightarrow \infty} n_j = \infty$. By H3, for j sufficiently large, the periodic points are hyperbolic. Therefore (y, λ_0) is not an explosion point. $\square$

Now consider the positive derivative case. Let w be a homoclinic tangency point contained in homoclinic orbit $(z_{-k})_{k=0}^\infty$. Since the eigenvalue of x_0 is positive, there is a M sufficiently large such that $(z_{-k})_{k=M}^\infty$ lies entirely on one side of x_0 . That is, the homoclinic orbit converges to the fixed point




Figure 4: The tangency point w is contained in $U_{left} \cap U_{right}$, and is thus not an explosion point. However, the preimages i and j of w are explosion points.

along one branch of the unstable manifold. We denote this by saying that the homoclinic orbit is contained in the local right or left branch of x_0 , as formalized in the following definition.

Definition 6 (Two branches of the unstable manifold). *Let x_0 be a repelling fixed point for $f \in C^2$, with $f'(x_0) > 0$. Locally, the left and right branches of the unstable manifold of x_0 are disjoint. Define U_{left} and U_{right} to be the respective unions of images of local left and right manifold branches.*

Remark 4. *The union of U_{left} and U_{right} is the entire unstable manifold of x_0 . By the intermediate value theorem U_{left} and U_{right} are intervals. If $(U_{left} \cap U_{right}) \setminus x_0$ is not empty, then the intersection must contain either U_{left} or U_{right} . For example, U_{right} may contain points both to the left and to the right of U_{left} . See Figure 4 and 5.*

From the proof of Theorem 3, it is clear that to study chain explosions in homoclinic orbits, it is sufficient to consider homoclinic orbits containing tangency points. We formalize the notation in the following hypothesis:

H6 For a family satisfying H1 and H4, at f_0 , point w is a homoclinic tangency point to x_0 contained in (at least one) homoclinic orbit $(z_{-k})_{k=0}^{\infty}$. Let L be such that $w = z_{-L}$.

Theorem 4 (No explosions when manifold branches intersect). *Assume H1-6. Assume that $f'_0(x_0) > 0$ and that $w \in U_{right}$. If for any neighborhood $N \ni w$, the sets $f_0^L(N)$ and $U_{right} \setminus x_0$ contain points in common, then (w, λ_0) is not an explosion point.*

Proof. The details of this proof are similar to the negative derivative case. Preimages of any small neighborhood N of y are contracting and in the local righthand manifold branch. Thus the L^{th} image of N includes a shrunk preimage of N . Therefore N contains a periodic point, which by H3 and H5 is hyperbolic. Since N is arbitrary, (w, λ_0) is not an explosion point. $\square$

Remark 5. *The theorem above is also true when U_{right} is replaced by U_{left} .*

Theorem 5. *Assume H1-6, and that w is contained in $U_{right} \cap U_{left}$. Then (w, λ_0) is not an explosion point.*

Proof. The image of a neighborhood of w under f_0^L contains either points of U_{right} or U_{left} . But w is contained in both U_{left} and U_{right} , so by Theorem 4, (w, λ_0) is not an explosion point. $\square$

As mentioned in the introduction, in our previous work we gave a useful geometric method of approaching chain explosions in homoclinic orbits in two and three dimensions, termed crossing cycles. In two dimensions, we showed that a crossing cycle is necessary and sufficient for a chain explosion to occur [1]. The analogous statements are true in one dimension, as in the theorem below. Crossing and non-crossing orbits are shown in Figures 5 and 2 respectively.

Definition 7 (Crossing orbits). *Assume H1-6. Note that for $g := f_0^{L-1}$, z_{-1} is a point of tangency. If for sufficiently large k , (i) $z_{-k} \in U_{left}$, and the graph of g is locally above the horizontal line at z_{-1} , or (ii) $z_{-k} \in U_{right}$, and the graph of g is locally below the horizontal line at z_{-1} , then we call the homoclinic orbit (z_{-k}) a **crossing orbit**. A homoclinic orbit that is not crossing is called a **non-crossing orbit**.*

Theorem 6 (Explosions imply crossing orbits). *If H1-6 hold and (w, λ_0) is an explosion point, then $(z_{-k})_{k=0}^\infty$ is a crossing orbit.*

Proof. If the graph of g is locally above the horizontal line at z_{-1} , then f_0^L of any neighborhood of w is contained in U_{right} . Likewise, if the graph is locally below the line, f_0^L of a neighborhood of w is contained in U_{left} . The result now follows from Theorem 4. $\square$




Figure 5: In this figure, w is a homoclinic tangency point contained in a crossing orbit. Thus w lies in U_{right} , but not U_{left} . By Theorem 7, w is an explosion point. As depicted here, the preimage z of w is also an explosion point.

The results stated so far give necessary conditions for an explosion point. We now give a converse to these. We first make a generic hypothesis.

By Theorem 5, to classify explosion points, it suffices to only consider the case of a tangency point w contained only in one manifold branch of x_0 , which we state in the following hypothesis.

H7 Assume H1-6. Assume that at λ_0 the tangency point w is contained exclusively in one manifold branch of x_0 .

Remark 6. Assume H1-7. Then at $\lambda = \lambda_0$, either tangency point $w \in U_{right}$, and U_{left} contains no tangencies to fixed points or periodic orbits, or tangency point $w \in U_{left}$, and U_{right} contains no tangencies to fixed points or periodic orbits. This follows from the fact that w is only contained in one of U_{left} and U_{right} , so by H3 there is no tangency in the other manifold branch.

The following theorem gives sufficient conditions for an explosion.

Theorem 7 (Crossing bifurcations and explosions). Assume H1-7. If every homoclinic orbit containing w is a crossing orbit, then (w, λ_0) is an explosion point.

Proof. For specificity, let $w \in U_{right}$. Since w is only contained in crossing orbits, w is not contained in U_{left} , and the L^{th} image of any small neighborhood of w is contained only in U_{left} . This implies that $U_{left} \subset U_{right}$, but that the subset is strict. As a result, prior to tangency, w is contained in U_{left} but not U_{right} . Thus the right endpoint of U_{left} is x_0 . The left endpoint x_L of U_{left} cannot map to x_0 , since that would imply a homoclinic tangency in U_{left} . In fact, no points in $U_{left} \setminus x_0$ map to x_0 , since there are assumed to be no tangencies and no intersections with U_{right} . If x_L maps into U_{left} , then there is a neighborhood of x_L which also maps into U_{left} . If x_L is a fixed point, then by H3 and H7, it is hyperbolic. It has no critical points to mapping to it from the interior of U_{left} . Thus x_L is an attracting fixed point. Prior to tangency, there is an $\epsilon > 0$ such that no ϵ -chains carry points from U_{left} to U_{right} , and thus from w to itself. Thus (w, λ_0) is an explosion point. $\square$

We are interested not only in explosion points which are themselves tangency points, but also in explosion points far from tangencies which are caused by tangencies. Theorem 3 contained such a result, but the other results were specifically about the tangency points. It is now straightforward to combine the previous results with Lemma 1 to get results for general homoclinic points. Since the chain recurrent set is invariant under forwards iteration, the image of non-crossing tangency point is a non-explosion point. However, there may be explosion points with iterates that are non-explosion points. For example, Figure 4 shows points of explosion which are not tangency points but are preimages of tangency points.

Theorem 8 (Explosions when manifold branches do not intersect). *Assume H1-6. If for all k , (z_{-k}, λ_0) is an explosion point, then the orbit $(z_{-k})_{k=0}^\infty$ is a crossing orbit.*

Proof. If the orbit is a non-crossing orbit, then there are hyperbolic periodic orbits limiting on every point in the orbit. $\square$

Theorem 9. *If H1-6 hold, and (w, λ_0) is an explosion point, then any pre-image z_{-k} in the homoclinic orbit of w is such that (z_{-k}, λ_0) is an explosion point.*

Proof. If (w, λ_0) is an explosion point, then by Lemma 1, z_{-k} is not chain recurrent for $\lambda < \lambda_0$. At λ_0 , z_{-k} is contained in a homoclinic orbit, and thus chain recurrent. $\square$

Remark 7. *This theorem only mentions preimages of w which are contained in a homoclinic orbit at λ_0 . There may be other preimages of w which are not contained in any homoclinic orbit, and are thus not chain recurrent at λ_0 .*

Theorem 10. *Assume H1-6. Assume that for $\lambda = \lambda_0$, $(z_{-k})_{k=0}^\infty$ is a crossing orbit, but w is also contained in some other homoclinic orbit $(\alpha_{-k})_{k=0}^\infty$ which is a non-crossing orbit. Thus (w, λ_0) is not an explosion point. Then the following statements hold:*

- i. For all $m > 0$, $(f^m(w), \lambda_0)$ is not an explosion point.*
- ii. Assume H6 and H7. If there exists $J > L$ such that z_{-J} is contained neither in $(\alpha_{-k})_{k=0}^\infty$ nor in any other non-crossing orbit, then for all $k > J$, (z_{-k}, λ_0) is an explosion point.*

Proof. The first statement holds since at λ_0 , any image of w is contained in $(\alpha_{-k})_{k=0}^\infty$, which is a non-crossing orbit. Thus by Theorem 8, this image is not an explosion point.

Let $J > L$ be as in Part *ii* of the theorem. For specificity, assume that $z_{-J} \in U_{right}$. Since z_{-J} is contained in a crossing orbit, it is not contained in U_{left} . A neighborhood of z_{-J} maps into U_{left} under f_0^J . If the tangency point w is in U_{right} , the rest of this proof is exactly the same as the proof of Theorem 7, and we conclude that (z_{-J}, λ_0) is an explosion point. Even if $w \in U_{left}$, if the forward orbit of w is in the interior of U_{left} , the proof follows as before. Furthermore the forward orbit of w cannot contain x_L , the left endpoint of U_{left} : Namely, since x_L is the endpoint of an invariant interval, this either implies a double homoclinic tangency in the forward orbit of w , which is ruled out by H3; or it implies both a tangency at w and a transverse homoclinic intersection at x_L in the orbit of w , which would mean that z_{-J} is contained in a non-crossing orbit. Finally, using the same proof as given for Theorem 9, for all $k > J$, (z_{-k}, λ_0) is an explosion point. $\square$

This completes the classification of when homoclinic points are explosion points.

Unlike the planar case, in one dimension the heteroclinic case reduces to the homoclinic case. That is, if there is a transverse heteroclinic cycle including an orbit from $\{p\}$ to $\{q\}$, which are hyperbolic periodic orbits, then both periodic orbits must be repellers. Further, the unstable manifold of $\{p\}$ contains the unstable manifold of $\{q\}$. Since we assume only one tangency, a heteroclinic tangency point is also a homoclinic tangency point.

4 General explosion classification

There are many previous results on the structure of ω -limit sets and chain recurrent sets for interval maps. Block and Coppel [5] showed that the chain recurrent set for maps of the interval can be classified as the set of points $\{x : x \in Q(x)\}$, where $Q(x)$ is the intersection of all asymptotically stable sets containing the limit set $\omega(x)$. They showed that $Q(x)$ is either an asymptotically stable periodic orbit, a set of asymptotically stable iterated intervals, or special type of set known as a solenoidal set. However, this classification is not as useful as it would appear, since $Q(x)$ is not the set of points in the chain component containing x . Block [4] also proved that an interval map has a homoclinic point if and only if it has a periodic point with period not a power of two. Block and Hart [6] improved on this result to show the existence of a homoclinic point to a given power of two implies a cascade of homoclinic bifurcations. Further, if a family of maps changes from zero to positive entropy, then there is a cascade of homoclinic bifurcations. Of most relevance to the topic of this paper are the works of Sarkovskii, Mañé [16], Blokh [7, 9], and Blokh, Bruckner, Humke, and Smital [8], where the detailed structure of all possible ω -limit sets is studied. The ω -limit set can be a Cantor set known as a basic set (Definition 9). Another interesting case occurs when the ω -limit set of a point is a limit of a period doubling cascade, known as a solenoid (Definition 10). See [3, 15] for a detailed characterization of solenoids. Blokh [9] showed that for C^2 -smooth maps, ω -limit sets are either periodic orbits, periodic transitive intervals, subsets of basic sets, or solenoids. We use this result to systematically show that for all possible explosions, there are saddle-node or tangency bifurcations.

The following definition describes points at which jumps in an ϵ -chain are required. We call these points barricades, as they serve to obstruct orbits. For example, at the bifurcation parameter, a saddle node point blocks the points on one side from reaching the other side.

Definition 8 (Barricade). *Assume H1 and H2. Let z be any point. Let $S_\epsilon = \omega(B_\epsilon(z))$. A point $y \in \lim_{\epsilon \rightarrow 0} S_\epsilon$ is called a **barricade** for z if it is blocking the orbit of z . That is, let $Z_\epsilon = \omega(B_\epsilon(y))$. Then $\lim_{\epsilon \rightarrow 0} Z_\epsilon$ contains points not contained in $\lim_{\epsilon \rightarrow 0} S_\epsilon$.*

We consider a point such that the ω -limit set is a fixed point or periodic orbit.

Theorem 11. *Assume H1-3. Assume that for f_0 , z is a point such that $\omega(z)$ is equal to a fixed point or periodic orbit $\{p\}$, and $\{p\}$ is a barricade. Then $\{p\}$ is either non-hyperbolic or is the image of a critical point.*

Proof. Assume that p is a fixed point, since otherwise we can let $g = f_0^n$. Since p is a barricade, $p \in S_\epsilon$, and $\lim_{\epsilon \rightarrow 0} Z_\epsilon \neq \lim_{\epsilon \rightarrow 0} S_\epsilon$ (using the notation from Definition 8). This implies that p cannot be an attracting fixed point, since for an attracting fixed point $\lim_{\epsilon \rightarrow 0} Z_\epsilon = p$. Thus if p is hyperbolic, it must be repelling, meaning that z is a preimage of p . Define K by $f^K(z) = p$. Since p is a barricade, for small ϵ , $f^K(B_\epsilon(z))$ is an interval on one side of p , implying that p is in the orbit of a critical point. $\square$

The previous theorem only indicates that a critical point exists. It still remains to be shown that the critical point is actually a homoclinic or heteroclinic point. We use the fact that all ω -limit sets for interval maps have been classified. The following theorem is useful in the proof of several subsequent theorems.

Theorem 12. *Assume H1-3, and that M is an invariant interval under f_0 , such that for all $\epsilon > 0$, there is an ϵ -chain from a point $x \in M$ to a point $y \notin M$. Then there is an ϵ -chain from an endpoint e of M to y , where e is fixed or period two. Furthermore, if x is not an endpoint of M , then either e is non-hyperbolic; or e is repelling, and there is an orbit of a critical point in M mapping onto e .*

Proof. Since $f(M) = M$, the only way for an ϵ -chain to exit M is through an ϵ -jump across one of the endpoints. Call the endpoint e . Thus there is a chain from e to y . Assume e does not map to itself or to the other endpoint of M . Then $f(e)$ is contained in the interior of M . But this means that no small ϵ -jump at e exits M , which is a contradiction. Thus e can be chosen to be either a fixed point or a period two point.

The orbit of e cannot be attracting, since then there would not be ϵ -chains from e to any other point. If the orbit of e is hyperbolic, then it is repelling, and there is a chain from x to e . If the orbit of x includes e , and x is not an endpoint of M , then the orbit of x contains a critical point, since M is invariant. If the orbit of x does not include e , then the limit set of x contains more than just a repelling periodic orbit. $\square$

The following result shows what happens if the backward limit set of a point contains a periodic orbit.

Theorem 13. *Assume H1-3. Let $\{p\}$ be a fixed point or periodic orbit for f_0 which is hyperbolic. Let z be a point such that for all $\epsilon > 0$ there is an ϵ -chain from $\{p\}$ to z , but z is not in the unstable manifold of $\{p\}$. Then there is a homoclinic tangency to a periodic orbit.*

Proof. Assume without loss of generality that p is a fixed point (since otherwise, we can use the same proof for an iterate of f_0). Since p is hyperbolic, it is repelling. The unstable manifold of p is an invariant interval, denoted $U(p)$. Since $z \notin U(p)$, there is a barricade point for p . By Theorem 12, there must be a barricade point which is a periodic endpoint of $U(p)$, with a critical point in $U(p)$ mapping to the endpoint. That is, there is a homoclinic tangency point for the periodic endpoint. $\square$

Combining Theorems 11 and 13, we conclude that if an explosion occurs at a point (z, λ_0) such that $\omega(z)$ is equal to a periodic orbit, then there is either a saddle-node bifurcation point or a tangency between stable and unstable manifolds. We now consider more general ω -limit sets. First consider the case where the ω -limit set of a point is an interval.

Theorem 14. *Assume H1-3, for f_0 , z is such that $\omega(z)$ is an interval M , and that there exists $x \in M$ such that e is a barricade for z . Then e is an endpoint of the interval, e is fixed or period two, and if e is hyperbolic, then there is a homoclinic tangency to e .*

Proof. Since $\omega(z) = M$, f_0 is transitive on M . Therefore the unstable manifold of any repelling periodic orbit in M contains all of M . A barricade must not be in the interior of $\omega(z)$. Thus e is an endpoint of M . The result now follows from Theorem 12. $\square$

We now consider the case of an ω -limit set that is contained in an invariant interval but is nowhere dense.

Definition 9 (Basic set). *Assume H1. Let $M = \cup_{k=1}^n I_k$ be an n -periodic cycle of intervals for a function f_0 . Define $B(M, f_0) = \{x \in M : \text{for every open interval } U \text{ of } x \text{ in } M, \overline{\text{orb}(U)} = M\}$. If $B(M, f_0)$ is infinite, then it is called a basic set.*

Theorem 15. *Assume H1, H2, and H3a. Assume that (z, λ_0) is an explosion point, and $\omega(z)$ is nowhere dense and is contained in a basic set $B(M, f_0)$. Then $\omega(z)$ is a periodic orbit.*

Proof. Assume that (z, λ_0) is an explosion point, and that $\omega(z) \subset B(M, f_0)$. If z is contained in an interval complementary to $B(M, f_0)$, then by Blokh [7], $\omega(z)$ is a periodic orbit.

Assume $\omega(z)$ is non-periodic, meaning z is contained in the basic set. By [9], $B(M, f_0)$ is contained in the closure of the periodic orbits for f_0 . Using H3a, there is a sequence of hyperbolic periodic points converging to z . Thus (z, λ_0) is not an explosion point. $\square$

By the above theorem combined with Theorems 11 and 13, if an explosion point has an ω -limit set which is a basic set, then there is either a saddle-node point or a tangency. The last possibility for an ω -limit set is a solenoid, as in the following definition.

Definition 10 (Solenoid). *Assume H1. Let $M_j = \cup_{k=1}^{n_j} I_k^j$ be a nested sequence of cycles of intervals for a function f_0 with least period n_j , $\lim_{j \rightarrow \infty} n_j = \infty$. Thus $f_0^{n_j}(I_1^j) = I_1^j$, n_j is increasing, and for each j , $I_1^{j+1} \subset I_1^j$. If the set $S = \cap_{j=1}^{\infty} M_j$ is nowhere dense, then S is called a solenoid or Feigenbaum-like set.*

Jiménez López has shown that solenoids are the boundary of chaos and order [15]. Blokh [9] demonstrated that solenoids and basic sets are disjoint. We prove the following result.

Theorem 16. *Assume H1, H2, and H3a. Assume that (z, λ_0) is an explosion point, and $\omega(z)$ is a solenoid S . Then there is an infinite sequence periodic orbits which are barricades with associated tangencies.*

Proof. Since solenoids and basic sets are closed, invariant, and disjoint, there is a neighborhood of solenoid S containing no basic sets. By definition, there is an infinite nested sequence of invariant cycles of intervals M_j containing S . Blokh [7] proved that the periodic orbits are dense in a neighborhood of S , meaning that z is not contained in S . Thus for all j sufficiently large, for all $\epsilon > 0$ there is an ϵ -chain from points in M_j to z , but z is not contained in M_j . By Theorem 12, there exists e_j , an endpoint of an interval the cycle of M_j which is periodic. By hypothesis H3a, for sufficiently large j , e_j is hyperbolic. Since for all $\epsilon > 0$ there is an ϵ -chain from e_j to z , e_j is a repeller for large j . There is an orbit of a critical point in M_j mapping onto e_j . Furthermore, z is not contained in the unstable manifold of e_j , since there is a nested sequence of invariant M_j not containing z . Theorem 13 implies that the critical point to e_j is a point of homoclinic tangency. $\square$

Remark 8. *If f_0 has a finite number of homoclinic and heteroclinic tangencies, as assumed in H3b, then the above theorem shows that there are no forward chains from solenoid S to a point outside of S .*

Remark 9. *We have shown that there is either a tangency or a non-hyperbolic critical point contained in the same chain component as z . Under a generic hypothesis (H3a), a non-hyperbolic periodic orbit is either codimension-one saddle-node or period doubling bifurcation. In fact, such an orbit is not a period doubling point, since the periodic orbit at a period-doubling bifurcation point is attracting.*

We now combine the results of this section to give a proof of the General explosion classification theorem.

Proof. (of the General explosion classification theorem) Assume that (z, λ_0) is an explosion point. The only possibilities for $\omega(z)$ are a periodic orbit, a cycle of intervals, a nowhere dense basic set, and a solenoid. Above, we have shown that in any of these cases, there is a periodic barricade point for z which is either non-hyperbolic, or there is a homoclinic or heteroclinic tangency. In fact, the case of a solenoid is ruled out by H3b. By Remark 9, a non-hyperbolic periodic orbit must necessarily be a saddle-node point. This completes the proof the theorem. $\square$

5 Acknowledgements

E.S. would like to thank Alexander Blokh and Victor Jiménez López for their helpful suggestions. J.A.Y. was partially supported by NSF Grant DMS0616585.

References

- [1] K. Alligood, E. Sander, and J. Yorke. Explosions: Global bifurcations at heteroclinic tangencies. *Ergodic Theory and Dynamical Systems*, 22(4):953–972, 2002.
- [2] K. Alligood, E. Sander, and J. Yorke. Three-dimensional crisis: Crossing bifurcations and unstable dimension variability. *Physical Review Letters*, 96(244103), 2006.

- [3] L. Alsedà, V. Jiménez López, and L. Snoha. All solenoids of piecewise smooth maps are period doubling. *Fund. Math.*, 157(2-3):121–138, 1998. Dedicated to the memory of Wiesław Szlenk.
- [4] L. Block. Homoclinic points of mappings of the interval. *Proc. Amer. Math. Soc.*, 72(3):576–580, 1978.
- [5] L. Block and W. Coppel. *Dynamics in One Dimension*, volume 1513 of *Lecture Notes in Mathematics*. Springer, Berlin, 1992.
- [6] L. Block and D. Hart. The bifurcation of homoclinic orbits of maps of the interval. *Ergodic Theory Dynamical Systems*, 2(2):131–138 (1983), 1982.
- [7] A. Blokh. Density of periodic orbits in ω -limit sets with the Hausdorff metric. *Real Anal. Exchange*, 24(2):503–521, 1998/99.
- [8] A. Blokh, A. Bruckner, P. Humke, and J. Smítal. The space of ω -limit sets of a continuous map of the interval. *Transactions of the American Mathematical Society*, 348(4):1357–1372, 1996.
- [9] A. M. Blokh. The “spectral” decomposition for one-dimensional maps. In *Dynamics reported*, volume 4 of *Dynam. Report. Expositions Dynam. Systems (N.S.)*, pages 1–59. Springer, Berlin, 1995.
- [10] C. Bonatti, L. Diaz, and M. Viana. *Dynamics beyond hyperbolicity*. Springer Verlag, Berlin, 2005.
- [11] R. Bowen. Markov partitions for axiom a diffeomorphisms. *American Journal of Mathematics*, 92:725–747, 1970.
- [12] C. Conley. *Isolated Invariant Sets and the Morse Index*. American Mathematical Society, Providence, R.I., 1978.
- [13] L. Díaz and J. Rocha. Heterodimensional cycles, partial hyperbolicity and limit dynamics. *Fundamenta Mathematicae*, 174(2):127–186, 2002.
- [14] V. Horita, N. Muiz, and P. R. Sabini. Non-periodic bifurcations of one-dimensional maps. *Ergodic Theory and Dynamical Systems*, Online, 2007.
- [15] V. Jiménez López. Period doubling is the boundary of chaos and of order in the C^1 -topology of interval maps. *Nonlinearity*, 15(3):817–839, 2002.
- [16] R. Mañé. Hyperbolicity, sinks and measure in one-dimensional dynamics. *Comm. Math. Phys.*, 100(4):495–524, 1985.
- [17] F. Marotto. Snap-back repellers imply chaos in R^n . *Journal of Mathematical Analysis and Applications*, 63:199–223, 1978.
- [18] S. Newhouse and J. Palis. Cycles and bifurcation theory. *Astérisque*, 31(44–140), 1976.
- [19] J. Palis and F. Takens. *Hyperbolicity & Sensitive Chaotic Dynamics at Homoclinic Bifurcations*. Cambridge University Press, Cambridge, 1993.

- [20] P. R. Sabini. *Non-periodic bifurcations at the boundary of hyperbolic systems*. PhD thesis, IMPA, 2001.
- [21] E. Sander. Homoclinic tangles for noninvertible maps. *Nonlinear Analysis*, 41(1-2):259–276, 2000.

AMS Subject Classification: 37.

E. Sander Department of Mathematical Sciences, George Mason University, 4400
University Dr., Fairfax, VA, 22030, USA.
E-mail: sander@math.gmu.edu

J.A. Yorke, IPST, University of Maryland, College Park, MD 20742, USA.
E-mail: yorke@ipst.umd.edu